




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61M 1/34, A61K 31/445, 38/19, 38/17</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/61085</b> <b>(43) International Publication Date:</b> 2 December 1999 (02.12.99)
<b>(21) International Application Number:</b> PCT/US99/11306 <b>(22) International Filing Date:</b> 21 May 1999 (21.05.99) <b>(30) Priority Data:</b> 09/083,307 22 May 1998 (22.05.98) US <b>(71)(72) Applicant and Inventor:</b> LENTZ, M., Rigdon [US/US]; 6347 Shadow Ridge Court, Brentwood, TN 37027 (US). <b>(74) Agents:</b> PABST, Patrea, L. et al.; Arnall Golden & Gregory, LLP, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).		<b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>

**(54) Title:** METHOD AND COMPOSITIONS FOR TREATMENT OF CANCERS**(57) Abstract**

A method to treat cancer uses ultrapheresis, refined to remove compounds of less than 120,000 daltons molecular weight, followed by administration of replacement fluid, to stimulate the patient's immune system to attack solid tumors. In the preferred embodiment, the patient is ultrafiltered using a capillary tube ultrafilter having a pore size of 0.02 to 0.05 microns, with a molecular weight cutoff of 120,000 daltons, sufficient to filter one blood volume. The preferred replacement fluid is ultrafiltered normal plasma.

The patient is preferably treated daily for three weeks, diagnostic tests conducted to verify that there has been shrinkage of the tumors, then the treatment regime is repeated. The treatment is preferably combined with an alternative therapy, for example, treatment with an anti-angiogenic compound, one or more cytokines such as TNF, gamma interferon, or IL-2, or a procoagulant compound. The treatment increases endogenous, local levels of cytokines, such as TNF. This provides a basis for an improved effect when combined with any treatment that enhances cytokine activity against the tumors, for example, treatments using alkylating agents, doxorubicin, carboplatinum, cisplatin, and taxol. Alternatively, the ultrapheresis treatment can be combined with local chemotherapy, systemic chemotherapy, and/or radiation.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

**METHOD AND COMPOSITIONS FOR TREATMENT OF CANCERS****Background of the Invention**

5           The present invention is generally in the field of enhancing an immune response, and particularly relates to the removal of inhibitors of immune mediators, in combination with anti-angiogenic compounds, cytokines, compounds inducing a procoagulant state, chemotherapeutics and/or radiation.

10           This application claims priority to U.S.S.N. 09/083,307 filed May 22, 1998.

          Conventional cancer therapy is based on the use of drugs and/or radiation which kills replicating cells, hopefully faster than the agents kill the patient's normal cells. Surgery is used to reduce tumor bulk, but has little  
15   impact once the cancer has metatized. Radiation is effective only in a localized area.

          The treatments can in themselves kill the patient, in the absence of maintenance therapy. For example, for some types of cancer, bone marrow transplants have been used to maintain the patient following treatment with  
20   otherwise fatal amounts of chemotherapy. Efficacy has not been proven for treatment of solid tumors, however. "Cocktails" of different chemotherapeutic agents and combinations of very high doses of chemotherapy with restorative agents, for example, granulocyte macrophage colony stimulating factor ("GM-CSF"), erythropoietin, thrombopoietin  
25   granulocyte stimulating factor, ("G-CSF"), macrophage colony stimulating factor ("M-CSF") and stem cell factor ("SCF") to restore platelet and white cell levels, have been used to treat aggressive cancers. Even with the supportive or restrictive therapy, side effects are severe.

          Other treatments have been tried in an attempt to improve mortality  
30   and morbidity. Vaccines to stimulate the patient's immune system have been attempted, but not with great success. Various cytokines, alone or in

combination, such as tumor necrosis factor, interferon gamma, and interleukin-2 ("IL-2") have been used to kill cancers, but have not produced cures. More recently, anti-angiogenic compounds such as thalidomide have been tried in compassionate use cases and shown to cause tumor remission.

5 In animal studies, compounds inducing a procoagulant state, such as an inhibitor of protein C, have been used to cause tumor remission. New studies have shown that inhibitors of cytokine receptors, such as tumor necrosis factor receptors ("TNF-Rs") which are released in a soluble form from tumor cells, in high concentrations relative to normal cells, may restore  
10 the immune system's attack on the tumor cells (Jablonska and Peitruska, Arch. Immunol. Ther. Exp. (Warsz) 1997, 45(5-6), 449-453; Chen, et al., J. Neuropathol. Exp. Neurol. 1997, 56(5), 541-550).

U.S. Patent No. 4,708,713 to Lentz describes an alternative method for treating cancer, involving ultrapheresis to remove compounds based on  
15 molecular weight, which promotes an immune attack on the tumors by the patient's own white cells.

Despite all of these efforts, many patients die from cancer; others are terribly mutilated. It is unlikely that any one therapy will be effective to cure all types of cancer.

20 It is therefore an object of the present invention to provide a method and compositions for treatment of solid tumors.

It is a further object of the present invention to provide a method and compositions that does not involve non-selective, extremely toxic, systemic chemotherapy.

### 25 **Summary of the Invention**

A method to treat cancer uses ultrapheresis, refined to remove compounds of less than 120,000 daltons molecular weight, followed by administration of replacement fluid, to stimulate the patient's immune system to attack solid tumors. In the preferred embodiment, the patient is  
30 ultrapheresed using a capillary tube ultrafilter or parallel plate filter having a molecular weight cutoff of 120,000 daltons, sufficient to filter at least one

blood volume. The preferred replacement fluid is ultrapheresed normal plasma. Alternatively, the patient is pheresed to selectively remove soluble receptor/inhibitors to soluble tissue necrosis factor receptor-1 ("sTNFR-1"), soluble tissue necrosis factor receptor-2 ("sTNFR-2"), soluble interleukin-2 receptor ("sIL-2R"), soluble interleukin-1 receptor ("sIL-1R"), soluble interleukin-6 receptor ("sIL-6R"), or soluble interferon-gamma receptor ("sIFN-gammaR"). These can be removed by binding to the cytokine, an epitope thereof, or an antibody to the receptor. These can be immobilized in the filter, in a column, or using other standard techniques for binding reactions to remove proteins from the blood or plasma of a patient. The patient is preferably treated daily for three weeks, diagnostic tests conducted to verify that there has been shrinkage of the tumors, then the treatment regime is repeated.

The treatment is preferably combined with an alternative therapy, for example, treatment with an anti-angiogenic compound, one or more cytokines such as TNF, gamma interferon, other interferons, or IL-2, or a procoagulant compound. The treatment increases the inflammation against tumors by allowing cytokines, such as TNF, to work effectively. This provides a basis for an improved effect when combined with any treatment that enhances cytokine activity against the tumors, for example, treatments using alkylating agents, doxorubicin, carboplatinum, cisplatinum, and taxol, and other drugs which may be synergistic in effect with "unblocked" cytokines. Alternatively, the ultrapheresis treatment can be combined with local chemotherapy, systemic chemotherapy, and/or radiation.

#### **Brief Description of the Drawings**

Figures 1 and 2 are schematics of the system for ultrapheresis.

#### **Detailed Description of the Invention**

The methods and devices disclosed herein are useful for treatment of patients with cancer, immune-mediated disorders, chronic parasitism, some viral diseases, and other disorders characterized by elevated levels of TNF receptors or inhibitors to IL-2, IL-6, gamma interferon, or other pro-

inflammatory signals as well as white cell activation. Examples demonstrate efficacy in treating cancer patients.

## **I. Ultrapheresis**

### **A. Ultrapheresis System**

#### **1. Filters**

5 The filter must be biocompatible, and suitable for contact with blood, without causing excessive activation of platelets or clotting. Devices will typically be either parallel plate filters or capillary membrane filters. These can be adapted from devices currently in use for kidney dialysis. The  
10 capillary membrane filters will typically have a surface area of between about 0.25 and 1 m<sup>2</sup> for use with children and between about 1 and 3 m<sup>2</sup> for use with adults. The parallel plate filters will typically have a surface area in the range from 0.1 and 2 cm<sup>2</sup>/ml of blood to be filtered.

The filter membranes will typically be a biocompatible or inert  
15 thermoplastic such as polycarbonate, polytetrafluorethylene (Teflon<sup>®</sup>), polypropylene, ethylene polyvinyl alcohol or polysulfone, having a pore size typically of between 0.02 and 0.05 microns in a capillary membrane filter and of between 0.04 and 0.08 microns in a parallel plate filter. The actual pore size that yields the desired cutoff of approximately 120,000 daltons is  
20 determined based on the fluid flow geometry, shear forces, flow rates, and surface area. The effective cutoff for a capillary membrane filter with a pore size of 0.03 microns is 120,000 daltons, with a sieving coefficient of between 10 and 30%. This results in only a trivial amount of IgG being removed from the patient's blood. The filter membrane should be less than about 25  
25 microns, preferably less than about 10 microns, thick. The permeable membrane should not cause blood clotting or otherwise react with the blood.

A preferred membrane is one in which the pores are made by electron beams directed perpendicularly to the surface because the size and density of the pores can be accurately controlled in this manner. The pores are  
30 essentially circular in cross section so the effective pore size is the actual pore size. The effective pore size of ultrafiltered media having pores with

non-circular cross sections shall be the diameter of a circular pore which will pass molecules or other components of an equivalent size to the molecules or other components which pass through the filter medium in question.

Suitable devices can be obtained from Asahi Chemical Company,  
5 Japan, and Kuraray Co., Ltd, 1-12-39, Umeda, ite-ku, Osaka 530, Japan.

Staged filters can also be used, which have different pore sizes and/or geometries or surfaces areas, to provide for a "staggered" removal of materials from the blood. Alternatively, although not at this time preferred, one can use differential centrifugation, to provide for an appropriate  
10 separation of blood components. Specific absorbing columns can also be employed to selectively remove specific cytokine and cellular inhibitors from the filtered plasma so that the ultrafiltrate (treated plasma) can be returned to the patient.

## 2. Process Controls and Fluid Handling

15 The patient will typically be connected to the blood processing device using standard intravenous tubing, with connections similar to those used for plateletpheresis, so that blood can be removed from the patient at one site and returned at another. The tubing is connected to a pump that controls the flow rate so that in the preferred embodiment one blood volume (based on  
20 approximately 7% of the total body weight) is processed over a period of approximately 2 ½ hours. The filtrate is then returned from the filtration device to the patient at the second site. Standard microprocessor controls can be used to regulate the blood flow, for example, by monitoring the volume of the blood products being removed, in combination with flow rate monitors  
25 and pump speed.

The entire system should first be flushed with saline and then treated with an anticoagulant or ant clotting agent, such as sodium heparin, to be sure that there are no locations within the system where blood clotting can occur. Moreover, small amounts of anticoagulants should be continually  
30 introduced into the blood stream directed to the ultrafilter to ensure than no clotting occurs during the filtration process. All of the surfaces of the system

which come in contact with the blood and fluids which are infused into the patient must be sterilized prior to commencing treatment.

Figure 1 illustrates a system for ultrapheresis. Blood is removed from a patient by means of a venous catheter 10 with the distal lead 11 thereof disposed in the superior vena cava 12 leading to the patient's heart 13. The blood passes through conduit 14 to a drip chamber 15 and then into pump 16 which controls the pressure of the blood to the separation unit 17, preferably an ultrafilter as shown, through conduit 18. A pressure gauge 19 is provided on conduit 14 to continually monitor arterial pressure. A syringe pump 20 feeds an anti-clotting drug such as sodium heparin to conduit 18 to prevent the clotting of blood in the ultrafilter 17. In the ultrafilter 17 the blood stream passes over the ultrafilter medium or membrane 21 under pressure. The blood fraction including the low molecular weight components passes through the membrane 21 and is discharged as permeate through conduit 22. The retentate or treated blood containing the high molecular weight components, which include whole blood cells and platelets, is discharged into conduit 23 which ultimately leads back to the patient. Volumetric pump 27 passes a controlled amount of permeate to a container 28 for containment and for measuring. Volumetric pump 30, which is preferably the same type and capacity as pump 27, pumps replacement fluid from a container 31 to conduit 32, which directs the fluid to conduit 23 where it mixes with the retentate or treated blood. The treated blood and other components are returned to the patient through venous catheter 34, the distal or discharge end of which is disposed in the brachiocephalic vein. The volumetric pumps 27 and 30 are preferably set either to pump the same total amount of fluid or to pump at the same rate, so that the same volume of fluid which is removed from the patient's blood stream as permeate is returned as replacement fluid. The blood stream in conduit 23 is passed through filter 36 to remove clots or other debris from the blood stream. A drip chamber 37 ensures that no significant quantities of air enter the patient's blood stream.



A pressure gauge 38 is provided to continually monitor venous blood pressure.

Figure 2 illustrates another embodiment wherein blood removed from a patient is first passed through conduit 30 to a first ultrafilter 31 to selectively separate a blood fraction with components having molecular weights less than about 1,000,000 Daltons. The retentate from this ultrafiltration which contains the high molecular weight components is returned through conduit 32 to the patient. The permeate from the first ultrafilter 30 is passed through conduit 33 to a second ultrafilter 34 where a blood fraction having a molecular weight below 30,000 is removed from the permeate stream from the first ultrafilter 30. The permeate from the second ultrafilter 34, which contains the very low molecular weight components such as salts and nutrients may be returned to the patient through conduit 38. The retentate from the second ultrafilter which contains blocking factors, IgG immunoglobulins and other components is discharged through conduit 36 and 13.

Blood should be pumped through the ultrafilter unit at sufficient pressure to cause the blood components having the immunosuppressive effects to pass through the filter but at a velocity which will not excessively shear or otherwise damage the blood cells passing over the membrane. Generally it has been found that the ratio of the area of the membrane to the amount of blood treated per hour should be within about 0.1 to 2 cm/mL. Differential pressure across the membrane should range from about 2 to 20 mM Hg.

### 25                    3.     Replacement Fluids

The patient must receive replacement fluids following filtration. The preferred replacement fluid is ultrapheresis normal plasma, for example, expired plasma obtained from the Red Cross, which has been filtered using the same filter as used to treat the patient. Alternatively, the patient can be administered normal albumin, or fresh frozen plasma diluted with saline.

## II. Treatment with Adjuvant Therapies

Standard ultrapheresis is conducted over a period of time until a positive indication is observed. This is typically based on diagnostic tests which show that there has been some reduction in tumor size or which suggests tumor inflammation. The patient is preferably treated daily for three weeks, diagnostic tests conducted to verify that there has been shrinkage of the tumors and/or inflammation, then the treatment regime is repeated.

Surgical (or vacuum) removal of necrotic material may be required prior to or during treatment to avoid toxicity associated with high tumor burden.

This procedure has been demonstrated to cause a significant response (greater than 50% reduction in size of tumors) in a variety of solid tumors in approximately 50% of patients who have failed all other treatment modalities. A tumor specific inflammatory response provoked by ultrapheresis has been documented in approximately 75% of patients in metastatic melanoma clinical trials. This inflammation is characterized by redness, swelling, warmth, and tenderness and is confined to tumors only. There has been no associated injury to non-cancerous tissue. This tumor specific inflammatory response has led to a 50% or greater reduction in the size of tumors in 50% of patients studied so far. Clinical trials have also demonstrated a 44% major reduction of tumor metastases in human breast cancer and prostate cancer.

Tumor specific inflammation has been observed in patients with metastatic colon cancer, ovarian cancer, lung cancer, head and neck cancer, cervical and endometrial cancers. In some cases, this inflammation has been followed by significant tumor regressions in each tumor type. The significance of response in such diverse tumor types strongly suggests that ultrapheresis modifies the patient's response to the tumor in favor of successful immunologic control of the tumor. Types of tumors that are

particularly sensitive to the ultrapheresis include epithelial tumors, sarcomas, melanomas and glioblastomas.

It would clearly be advantageous to cause complete remissions. Based on the presumed mechanism that the process is removing immune inhibitors produced by the tumors, especially inhibitors of cytokines and other immune mediators, it is possible to treat the patients with adjuvant or combination therapies, that enhance the results achieved with the ultrapheresis. TNF-alpha and beta receptors are thought to be particularly important immune inhibitors. Therefore, compounds which enhance TNF activity are particularly preferred. These include anti-angiogenic compounds, such as thalidomide, procoagulant compounds, cytokines and other immunostimulants. Standard chemotherapeutic agents and/or radiation can also be used with the ultrapheresis.

#### A. Anti-Angiogenic Compounds

Any anti-angiogenic compound can be used. Exemplary anti-angiogenic compounds include O-substituted fumagillol and derivatives thereof, such as TNP-470, described in U.S. Patent No. 5,135,919, 5,698,586, and 5,290,807 to Kishimoto, et al.; angiostatin and endostatin, described in U.S. Patent No. 5,290,807, 5,639,725 and 5,733,876 to O'Reilly; thalidomide, as described in U.S. Patent Nos. 5,629,327 and 5,712,291 to D'Amato; and other compounds, such as the anti-invasive factor, retinoic acid, and paclitaxel, described in U.S. Patent No. 5,716,981 to Hunter, et al., and the metalloproteinase inhibitors described in U.S. Patent No. 5,713,491 to Murphy, et al. Thalidomide is administered once daily, 200 mg orally.

#### B. Procoagulant Compounds

Protein C is a vitamin K-dependent plasma protein zymogen to a serine protease. Upon activation it becomes a potent anticoagulant. Activated protein C acts through the specific proteolysis of the procoagulant cofactors, factor VIIIa and factor Va. This activity requires the presence of another vitamin K-dependent protein, protein S, calcium and a phospholipid (presumably cellular) surface. As described in Hemostasis and Thrombosis.

- Basic Principles and Clinical Practice 2nd Ed., Colman, R.W., et al., p. 263 (J.B. Lippincott, Philadelphia, PA 1987), protein C circulates in a two-chain form, with the larger, heavy chain bound to the smaller light chain through a single disulfide link. Protein C is activated to activated protein C (APC).
- 5 Thrombin is capable of activating protein C by the specific cleavage of the Arg<sup>12</sup>-Leu<sup>13</sup> bond in the heavy chain. *In vivo*, in the presence of physiological concentrations of calcium, the rate of this activation is enhanced dramatically when thrombin is bound to the endothelial cell cofactor, thrombomodulin. Matschiner, et al., Current Advances in Vitamin
- 10 K Research, pp. 135-140, John W. Suttie, ed. (Elsevier Science Publishing Co., Inc. 1988) have further reviewed the role of the Vitamin K dependent proteins in coagulation.

- Blockage of the natural anticoagulant pathways, in particular the protein C pathway, uses the natural procoagulant properties of the tumor to
- 15 target the tumor capillaries for microvascular thrombosis, leading to hemorrhagic necrosis of the tumor, as described in U.S. Patent No. 5,147,638 to Esmon, et al. Examples of such compounds include anti-protein C and anti-protein S.

- C. Cytokines** The biologic activity and clinical effectiveness
- 20 of pro-inflammatory cytokines is augmented by ultrapheresis in the patient with cancer and other states of acquired immune tolerance. Specifically, both TNF alpha and TNF beta, in doses of between approximately 100 to 500 micrograms per meter squared body surface area (M2BSA), can enhance the immune reaction in aggressive tumors. Monocyte and lymphocyte
- 25 activation is augmented by INF-alpha, INF-beta and gamma. The IL-1 and IL-2 receptor antagonists are removed by ultrapheresis and thereby upregulate the *in vivo* activity of these cytokines. An 80 kD glycoprotein, which is responsible for inhibiting blastoid transformation in advanced malignancy, chronic infectious disease and pregnancy, has recently been
- 30 found, and appears to be responsible for the loss of delayed hypersensitivity reactions in these diseases, which is removed by this process. This is

significant because in removing this type of suppression, vaccines of all types will work better. Dosage regimes for IFN-alpha and beta are 3 M units subcutaneous three times a week up to 20 M units/M2 BSA daily.

Interferon-gamma is administered in a dosage of between 100 to 1000 micgms per day.

#### **D. Anti-cytokine receptor molecules.**

It is well established that TNF receptor 1 and TNF receptor 2 molecules are shed by tumor cells, and that these molecules appear to inhibit immune mediated attack by the host on the tumor cells. The ultrapheresis is believed to remove the majority of these soluble receptors. Additional, and/or selective, removal of these soluble receptor/inhibitors to soluble tissue necrosis factor receptor-1 ("sTNFR-1"), soluble tissue necrosis factor receptor-2 ("sTNFR-2"), soluble interleukin-2 receptor ("sIL-2R"), soluble interleukin-1 receptor ("sIL-1R"), soluble interleukin-6 receptor ("sIL-6R"), or soluble interferon-gamma receptor ("sIFN-gammaR") can be used to supplement or instead of the ultrapheresis. The advantage of selective removal is that the same beneficial effect is obtained in treatment of the disorder but the treatment is much less expensive and safer since exogenous plasma or albumin does not have to be administered to the patient when there is selective removal. These can be removed by binding to the cytokine, an epitope thereof, or an antibody to the receptor. These can be immobilized in the filter, in a column, or using other standard techniques for binding reactions to remove proteins from the blood or plasma of a patient. As used herein, antibody refers to antibody, or antibody fragments (single chain, recombinant, or humanized), immunoreactive against the receptor molecules. In the preferred embodiment, these antibodies are immobilized on the ultrapheresis membrane filters, using standard antibody coupling techniques. In the most preferred embodiment, the antibody is reactive with the carboxy-terminus of the shed receptor molecules, thereby avoid concerns with signal transduction by the receptor is still present on the cell surface.

### E. Chemotherapeutic Agents

Preferred chemotherapeutic agents are those which are synergistic with TNF, for example, alkylating agents, doxyrubicin, carboplatinum, cisplatinum, and tomoxifen. Tamoxifen plays a role not only in blocking of estrogen receptors but also certain growth factor receptors such as epidermal derived growth factor ("EDGF"), fibroblast derived growth factor ("FDGF"), tumor derived growth factor ("TDGF"), TDGF- $\beta$  and platelet derived growth factor ("PDGF") and therefore may be complementary to inflammation against cancers provoked by ultrapheresis.

### F. Radiation

Radiation therapy is destructive of normal tissue, causing tumors to die partially by an inflammatory attack. Ultrapheresis allows the use of lower doses of radiation to kill residual tumor cells and spare normal tissue. In a preferred method, ultrapheresis is used as the initial therapy, followed by radiation at approximately one-half of the normal dosages. It is well established that TNF kills tumor cells by generating free oxygen radicals, hydroxyl radicals and halide ions, and that radiation therapy generates carbonium ions in tissue. Therefore the combination of the two is more effective in killing cancer cells than either alone.

### III. Examples

#### Example 1: Treatment of A Patient with Metastatic Leiomyosarcoma with Ultrapheresis.

Mrs. J.K. is a 43 year old lady with metastatic leiomyosarcoma with six (6) lung metastases all of which developed within one month of surgery on both lungs to remove tumors. These tumors had also failed methotrexate, adriamycin, ifosphomide, and dactinomycin.

She received 24 ultrapheresis procedures with no side effects. One month later, CAT scan revealed only four (4) tumors and these were reduced in size by 50%.

**Example 2: Treatment of Patients with Breast Cancer by  
Ultrapheresis and Thalidomide.**

Mrs. J.R. is a 44 year old lady who had metastatic breast cancer that had failed radiation therapy and treatment with chemotherapeutic agents: cytoxan, adriamycin, 5-FU, taxol, cis-platin, navalbine, tamoxofin and arimedx. Tumor at the time of ultrapheresis was documented in lungs, bone and skin of the entire left anterior and lateral chest.

She was treated with 15 ultrapheresis procedures over a three week period. She experienced marked inflammation in the tumors of his skin, increased pain from the tumors in her bones, and swelling of the tumors in her lungs. She then received the drug thalidomide 200 mg at night. The redness and swelling in her skin improved within 2 days and her breathing returned to normal within one week. Two weeks after completing treatment, all tumor in her skin had resolved clinically, her bone pain resolved and the tumors in her lungs resolved on repeat CAT scan. One week later, she returned to work as a school counselor. She tested disease free two months after treatment and was being maintained on thalidomide at the same dose.

**Example 3: Treatment of Patient with Metastatic Melanoma with  
Ultrapheresis and Thalidomide.**

Mr. P.G. is a 54 year old engineer with metastatic melanoma with metastases to lung and to lymph nodes in the mediastinum.

He received 24 ultrapheresis procedures, resulting in a 25% reduction of tumors. He was subsequently treated with an additional 12 procedures, resulting in minor tumor reduction despite evidence of tumor inflammation. The tumors regrew within one month. He was again retreated with ultrapheresis, again resulting in inflammation and some minor regression, but was then treated with thalidomide at the time of tumor inflammation. Two months later, repeat CAT scan showed complete disappearance of tumors in the lung and mediastinum. He is being followed closely and shows no evidence of disease and has no medical complaints six months after completing treatment.

**Example 4: Treatment of a Patient with Metastatic Adenocarcinoma  
with Ultrapheresis and Thalidomide.**

Dr. R.S. is a 59 year old gentlemen with metastatic adenocarcinoma of the left upper lung with metastases to liver, brain and bones. His tumors  
5 had failed to respond to taxol, cis-platin and etoposide. His brain tumors had responded to radiation therapy.

He received 15 ultrapheresis procedures. Each procedure caused increased pain in tumors of his spine, pelvis, right hip and left shoulder. Follow up scans after ultrapheresis treatment revealed resolution of tumors in  
10 pelvis, spine, hip, and ribs. There was a 50% reduction in the primary tumor in the lung and liver. Thalidomide was then started at 200 mg each night. One month later, the scans revealed further reduction in the tumors in lung and liver. The patient's pains have all been resolved and he is asymptomatic at this time.



22. A kit for treatment of a patient to induce an immune response against transformed, infected or diseased tissue comprising:

a device for removing only components present in the blood having a molecular weight of 120,000 daltons or less and

an agent selected from the group consisting of anti-angiogenic compounds, procoagulant compounds, cytokines, chemotherapeutic agents, and radiation, in a dosage formulation for treatment of the patient.

23. The kit of claim 22 wherein the agent is an anti-angiogenic compound.

24. The kit of claim 22 wherein the agent is a procoagulant compound.

25. The kit of claim 22 wherein the agent is a cytokine.

26. The kit of claim 25 wherein the cytokine is selected from the group consisting of GM-CSF, erythropoietin, thrombopoietin, G-CSF, M-CSF and SCF.

27. The kit of claim 22 wherein the agent is a chemotherapeutic agent.

28. The kit of claim 27 wherein the agent is selected from the group consisting of alkylating agents, doxorubicin, carboplatinum, cisplatinum, and taxol.

29. The kit of claim 22 further comprising anticoagulant to treat the device for removal of components from the blood prior to use.




Fig. 1




Fig. 2



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61M 1/34, A61K 31/445, 38/19, 38/17</b>		<b>A3</b>	(11) International Publication Number: <b>WO 99/61085</b>
			(43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11306		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 21 May 1999 (21.05.99)			
(30) Priority Data: 09/083,307 22 May 1998 (22.05.98) US		Published With international search report.	
(71)(72) Applicant and Inventor: LENTZ, M., Rigdon [US/US]; 6347 Shadow Ridge Court, Brentwood, TN 37027 (US).		(88) Date of publication of the international search report: 23 March 2000 (23.03.00)	
(74) Agents: PABST, Patrea, L. et al.; Arnall Golden & Gregory, LLP, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).			

(54) Title: METHOD AND COMPOSITIONS FOR TREATMENT OF CANCERS

## (57) Abstract

A method to treat cancer uses ultrapheresis, refined to remove compounds of less than 120,000 daltons molecular weight, followed by administration of replacement fluid, to stimulate the patient's immune system to attack solid tumors. In the preferred embodiment, the patient is ultrapheresed using a capillary tube ultrafilter having a pore size of 0.02 to 0.05 microns, with a molecular weight cutoff of 120,000 daltons, sufficient to filter one blood volume. The preferred replacement fluid is ultrapheresed normal plasma. The patient is preferably treated daily for three weeks, diagnostic tests conducted to verify that there has been shrinkage of the tumors, then the treatment regime is repeated. The treatment is preferably combined with an alternative therapy, for example, treatment with an anti-angiogenic compound, one or more cytokines such as TNF, gamma interferon, or IL-2, or a procoagulant compound. The treatment increases endogenous, local levels of cytokines, such as TNF. This provides a basis for an improved effect when combined with any treatment that enhances cytokine activity against the tumors, for example, treatments using alkylating agents, doxorubicin, carboplatinum, cisplatinum, and taxol. Alternatively, the ultrapheresis treatment can be combined with local chemotherapy, systemic chemotherapy, and/or radiation.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/11306

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61M1/34 A61K31/445 A61K38/19 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61M A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 708 713 A (LENTZ M.R.) 24 November 1987 (1987-11-24)  claims 1-13	1-5, 12-15, 20-22
A	US 5 679 260 A (BOOS K. ET AL.) 21 October 1997 (1997-10-21) the whole document	1-29
A	DE 196 24 250 A (AHRENHOLZ P. ET AL.) 2 January 1998 (1998-01-02) the whole document	1-29
A	US 4 486 282 A (BIER M.) 4 December 1984 (1984-12-04) the whole document	1-29
	--- -/-- ---	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

22 December 1999

Date of mailing of the international search report

12/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 apo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Moreau, J

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/11306

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DE 43 45 200 A (FRESENIUS AG)  22 December 1994 (1994-12-22)</p> <p>-----</p>	1-29

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/11306

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
**Remark: Although claims 1-11 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

information on patent family members

Initial Application No

PCT/US 99/11306

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4708713 A	24-11-1987	AT 88097 T	15-04-1993
		AU 596567 B	10-05-1990
		AU 4992985 A	22-05-1986
		CA 1262681 A	07-11-1989
		DE 3587266 A	19-05-1993
		EP 0184040 A	11-06-1986
		JP 61172563 A	04-08-1986
		MX 168132 B	06-05-1993
US 5679260 A	21-10-1997	DE 4435612 A	11-04-1996
		EP 0705845 A	10-04-1996
		JP 2807197 B	08-10-1998
		JP 8193031 A	30-07-1996
DE 19624250 A	02-01-1998	NONE	
US 4486282 A	04-12-1984	NONE	
DE 4345200 A	22-12-1994	DE 4312916 A	20-10-1994
		AU 6677894 A	08-11-1994
		WO 9423747 A	27-10-1994